

THE 7TH INTERNATIONAL CONFERENCE ON FOOD AND APPLIED BIOSCIENCE 2024

🕒 February 8 - 9, 2024

BOOK OF ABSTRACT FAB 2024

“ FOOD AND BIOTECHNOLOGY
FOR A SUSTAINABLE & BIO-CIRCULAR-GREEN ECONOMY ”



International Conference on Food and Applied Bioscience

February 8-9, 2024

Kantary Hills Hotel, Chiang Mai

BOOK OF ABSTRACTS

Organizer



Co-Hosts



The International Conference on Food and Applied Bioscience 2024
February 8-9, 2024 Kantary Hills Hotel, Chiang Mai, Thailand

Conference Committee

Academic Advisory Committee:

Assistant Professor Dr. Sujinda Sriwattana Chiang Mai University

Administrative Chairperson:

Associate Professor Dr. Pichaya Poonlarp Chiang Mai University

Editorial board/Scientific Committee

Professor Dr. Alissara Reungsang	Khon Kaen University
Professor Dr. Charles Brennan	STEM College
Professor Dr. Kyu Jin Park	Korea University
Professor Dr. Lee Youn Suk	Yonsei University
Professor Dr. Masanori Watanabe	Yamagata University
Professor Dr. Pornchai Rachtanapun	Chiang Mai University
Associate Professor Dr. Phisit Seesuriyachan	Chiang Mai University
Associate Professor Dr. Piyaporn Chueamchaitrakun	Mae Fah Luang University
Associate Professor Dr. Saowapa Chaiwong	Mae Fah Luang University
Associate Professor Dr. Tuyen Chan Kha	Nong Lam University
Associate Professor Dr. Yuthana Phimolsiripol	Chiang Mai University
Assistant Professor Dr. Toya Ishii	Kagawa University
Dr. Nguyeu Minh Xuan Hong	Nong Lam University

Organizer

Faculty of Agro-Industry, Chiang Mai University

Co-hosts

College of Maritime Studies and Management, Chiang Mai University

Food Science & Technology Association of Thailand

Korea University

Yamagata University

Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences

Agro-Industry Academic Council Association

Nong Lam University

Yonsei University

Guangzhou Institute of Energy Conversion, Chinese Academy of Sciences

Universitas Brawijaya

**P - 022 Phenolic Profile of Serbian Old Wheat Local Landraces
and Locally Adapted Varieties**

Aleksandra Mišan^{1,*}, Bojana Radić¹, Nikola Maravić¹, Milica Pojić¹,
Jelena Tomić¹, Miroslav Hadnađev¹ and Sanja Mikić²

¹*Institute of Food Technology, University of Novi Sad, Novi Sad, Serbia*

²*Institute of Field and Vegetable Crops, National Institute of the Republic of Serbia, Novi Sad, Serbia*

**Corresponding author E-mail: aleksandra.misan@fins.uns.ac.rs*

Abstract

Wheat is one of the most important staple crops in Serbia and plays an important role in food security and nutrition. Phenolic compounds are dominant antioxidants in whole grains and are essential quality traits in future breeding programs as they are known for their beneficial effects on health. With the aim of analyzing local wheat diversity in terms of phenolic compounds, 33 local landraces and old locally adapted varieties and 3 modern wheat varieties were grown in 2021/2022. So, the objective of this work was to optimize the extraction of phenolic compounds and analyze them by HPLC/DAD method and compare old with modern cultivars. As it is known that cereal grains contain different phenolic compounds, mainly phenolic acids in bound form, the extraction of phenolic compounds was done from wholemeal flour samples after hydrolysis. Alkaline hydrolysis conducted at 25°C for 12 h using a 2 M NaOH solution in water demonstrated greater efficiency compared to employing solutions of up to 10 M KOH in methanol under the same temperature and duration. HPLC-DAD analysis revealed that ferulic acid presents the major phenolic compound both in old and modern wheat samples ranging between 14 and 149 mg/kg in old and 51 and 61 mg/kg in modern varieties. Apart from ferulic acid, vanillic acid (3.1-11.5 mg/kg), syringic acid (1.4-5.5 mg/kg) and *p*-coumaric acid (1-6.9 mg/kg) were identified and quantified in old, but not in new varieties. Our results indicate that the genetic potential of old varieties in terms of polyphenolic compounds is not fully utilized because the content of ferulic acid in whole grains in some samples is almost three times higher than in modern varieties that do not differ significantly among themselves.

Keywords: Wheat; Phenolic profile; Ferulic acid; HPLC